

Mucopyocele of the Maxillary Sinus: A Case Study

Maha Hakimi, ENT - Neck and Head surgery Department; University medical center Mohammed VI. Marrakech
 Hadid Fatima Zahra, ENT - Neck and Head surgery Department; University medical center Mohammed VI. Marrakech
 Lakhdar Youssef, ENT - Neck and Head surgery Department; University medical center Mohammed VI. Marrakech
 Benhoummad Othmane, ENT- Neck and Head surgery Department; University medical center Mohammed VI. Marrakech
 Youssef Rochdi, ENT - Neck and Head surgery Department; University medical center Mohammed VI. Marrakech
 Abdelaziz Raji, ENT - Neck and Head surgery Department; University medical center Mohammed VI. Marrakech

Abstract:

Introduction: Mucocele is a benign but expansive cystic formation, lined by a respiratory epithelium. When its content is infected, it is called a mucopyocele. It usually develops in the frontal-ethmoid complex. The maxillary sinus location is exceptional.

Case report: a 43-year-old man, presented with chronic unilateral nasal obstruction, purulent rhinorrhea and anosmia. Rhinoscopy showed a bulge in the internal wall of the maxillary sinus emanating from the medium meatus. The computed tomography (CT) showed complete filling of the left maxillary with low-density mass. Magnetic resonance imaging (MRI) confirmed the diagnosis of mucopyocele of the left maxillary sinus. Patient underwent endoscopic surgery with complete recovery.

Discussion/conclusion: Mucopyocele of the maxillary sinus is a benign rare lesion, however destructive. A radiological assessment is essential to guide the choice of surgical treatment. The endonasal route remains the reference treatment for this pathology.

Keywords:- Maxillary Sinus, Mucopyocele, Mucocele.

I. INTRODUCTION

Mucocele is defined as a mucus-filled cavity that can occur in the oral cavity, appendix, gallbladder, paranasal sinuses or lacrimal sac. [1] When it is secondarily infected, as a natural though rare evolution of the illness it is called a mucopyocele. The maxillary sinus localization is quite rare, by far overridden by the frontal and anterior ethmoid sinuses localization. [2] In this article we described a rare case of mucopyocele of the maxillary sinus in an adult and dived into its pathology, treatment and outcome.

II. CASE REPORT

A 43-year-old male with no relevant past medical, surgical or traumatic history, reported with a chief complaint of left purulent rhinorrhea, unilateral nasal obstruction and anosmia evolving for 8 months. There was no history of any mass in the facial region, or any ocular or dental complains, or any paresthesia or numbness present in the region of chief complaint. Moreover, the patient complained of two episodes of epistaxis. The extra-oral examination showed slight

tenderness in the maxillary region without any evidence of swelling or inflammation. Well as, the intra oral examination showed a poor dental status with no swelling or inflammation. Rhinoscopy findings included an inflammatory state of the mucosa with a bulge in the internal wall of the maxillary sinus emanating from the medium meatus. The ophthalmological examination was normal.

The CT showed complete obliteration of the left maxillary sinus with soft-tissue density mass, responsible of a thinning and bulging of the inner wall of the maxillary sinus suggestive of benign expansive and destructive lesion of the left maxillary sinus, it showed a filling of the ostium and the left nasal cavity. In order to properly diagnose and to judge the extent of the pathology an MRI was advised, showing a mass filling the totality of the left maxillary sinus with a T1 hyper signal and associated with a T2 hyper signal. Moreover, it was achieving a mass effect on the wall separating the maxillary sinus and the nasal cavity, exteriorized in the nasal cavity until it made contact with the nasal septum deviating it to the right, and with no effect on the nasal floor. Which was in favor of a mucopyocele of the maxillary sinus. There was no intracerebral or intraorbital extension and all the other sinuses were unscathed (Figure 1 ,2). The blood work was normal, including a normal blood cell count and a normal C reactive protein rate.



Figure 1: complete obliteration of the left maxillary sinus with soft-tissue density mass



Figure 2: mass filling the totality of the left maxillary sinus

The patient underwent an incisional biopsy of the lesion under local anesthesia and the histopathological report was suggestive of infected mucocele of maxillary sinus (mucopyocele). The patient underwent a medical preparation in order to go through surgery consisting in antibiotics and steroids treatment. A surgical excision of the entire lesion was carried out under general anesthesia, and he benefited from a large middle meatal antrostomy (marsupialization). The histopathological examination afterwards was coherent with the diagnosis of mucopyocele. Patient recovery was uneventful up to one year after the procedure with no fresh complaints. (Figure 3)



Figure 3: Post-operative aspect of the antrostomy

III. DISCUSSION

Mucoceles and mucopyoceles are rare cystic and cavitated collections originating in the paranasal sinuses. They most commonly occur in the frontal and ethmoid sinuses; therefore, the maxillary sinus localization is quite rare. A mucocele contains mucinous secretions, whereas a mucopyocele is an abscess-like formation that also contains purulent material. [3]

Their origin has been the subject of much controversy but several theories have been proposed to try to explain the origin of this benign but destructive pathology. Mainly, the infectious theory secondary to the obstruction of the sinuses ostium, the inflammatory theory or the hypersecretory theory where the release of bone resorption factors causes the disease and finally the traumatic theory. The vicious circle: inflammation and ostial obstruction summarizes the cause of mucopyocele. [4] In this case, we can consider the bad dental state as a precipitating factor for the development of mucocele, it is after all an inflammatory nest. The incidence of mucocele in the general population is 0.4-0.8% [5], and the majority of mucocele occur in age between 40 to 70 years. There is a slight male predominance and right and left sides are equally affected. [6] These results clearly concur with our case, a 43-year-old male.

The usual symptoms of mucopyocele of the maxillary sinus include nasal obstruction, cheek pressure or pain, rhinorrhea and anosmia [7] These symptoms may be associated with ophthalmological or neurological signs such as headaches, orbital pain or exophthalmos. Cases of dacryocystocele, intra-orbital extension, and subdural empyema have been reported in the literature [8, 9].

The diagnosis has greatly benefited from the contribution of imaging especially paranasal sinuses CT. On computed tomography (CT) the mucocele is an opacity that is often regular, expansive, blowing the bone walls and pushing back more or less neighboring structures. Magnetic resonance Imaging (MRI), also, allows to perfectly appreciate the extension of the mucocele and to specify at most its relationship with adjacent structures. It also permits the diagnosis by showing a typical appearance of a cystic pocket with well-defined walls. The signal is variable depending on the viscosity and protein content of the intramucosal retention. So, it usually presents a T1 hypersignal associated with a variable T2 signal (hyper- and hyposignal depending on the protein concentration in the cavity). [10] A microbiological study of 36 mucopyoceles from different sinuses and not exclusively from the maxillary sinus showed a polymicrobial aerobic and anaerobic bacteriology, such as staphylococcus aureus, alpha-hemolytic streptococci, Hemophilus spp, Gram-negative bacilli, Peptostreptococcus sp., Prevotella sp., Fusobacterium sp., and Propionibacterium acnes [11].

An early management is of the most importance aspect of the treatment in order to prevent any orbital and intracranial complication. The recommended treatment of this entity is based on the endonasal endoscopic surgery. In general, this approach is preferred in paranasal sinus mucoceles due to its many advantages including a minimally invasive surgery and a lesser recurrence rate (4.8%) [12]. The principle of this surgery is to widen up the mucopyocele so to evacuate its content, and to create a large communication between the mucocele pocket and the nasal cavity (1). Different methods are used: lateral marsupialization, medial marsupialization or a transverse excision. Here, an incision of the pocket was made and supplemented by the opening of the cavity in order to prevent recurrence.

IV. CONCLUSION

Though it is a rare entity, the diagnosis, management and treatment of the mucopyocele of the maxillary sinus should be in the gear of every practitioner, especially due to the intra cerebral complications it can cause which are life threatening.

Conflict of interest

None

Funding

None

Consent

Informed consent was obtained from the patient for publication of this case report and accompanying images.

REFERENCES

- [1]. Baurmash HD. Mucoceles and ranulas. J Oral Maxillofac Surg. 2003;61:369–78.
- [2]. Avinash Kshar, Abhijeet Patil, Hemant Umarji, and Sonali Kadam Mucopyocele of the maxillary sinus ; 2014 Jan-Feb; 11(1): 119–123
- [3]. Gabriel Zada M. , Beatriz S. Lopes; Mucocele and mucopyocele, Atlas of Sellar and Parasellar Lesions pp 411-414.
- [4]. Bahadir O, Imamoglu M, Bektas D. Massive concha bullosa pyocele with orbital extention. Auris Nasus Larynx 2006 ; 33 : 195–198..
- [5]. Ozturk K, Yaman H, Arbag H, Koroglu D, Toy H. Submandibular gland mucocele: Report of two cases. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2005;100:732–5.
- [6]. Kennedy DW, Josephson JS, Zinreich SJ, Mattox DE, Goldsmith MM. Endoscopic sinus surgery for mucoceles: a viable alternative. Laryngoscope. 1989;99:885–889
- [7]. Busaba NY1, Salman SD. Maxillary sinus mucoceles: clinical presentation and long-term results of endoscopic surgical treatment 1999 Sep;109(9):1446-9.
- [8]. Barry J. Edison, Dale R. Meyer. Nasolacrimal Duct Obstruction and Dacryocystocele Associated with a Concha Bullosa Mucocele. Ophthalmology 2000;107:1393–1396